



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification<sup>6</sup> : C12N 15/12, 1/21, A61K 38/38, C12P 21/02, C12N 15/11, 1/20</p>	<p>A1</p>	<p>(11) International Publication Number: <b>WO 95/23857</b> (43) International Publication Date: 8 September 1995 (08.09.95)</p>
<p>(21) International Application Number: PCT/GB95/00434 (22) International Filing Date: 1 March 1995 (01.03.95) (30) Priority Data: 9404270.2 5 March 1994 (05.03.94) GB (71) Applicant (for all designated States except US): DELTA BIOTECHNOLOGY LIMITED [GB/GB]; Castle Court, Castle Boulevard, Nottingham NG7 1FD (GB). (72) Inventors; and (75) Inventors/Applicants (for US only): KERRY-WILLIAMS, Sean, Martin [GB/GB]; 131 Trent Road, Beeston Rylands, Nottingham NG9 1LP (GB). GILBERT, Sarah, Catherine [GB/GB]; 65 Dene Road, Headington, Oxford OX3 7EQ (GB). (74) Agents: BASSETT, Richard et al.; Eric Potter Clarkson, St. Mary's Court, St. Mary's Gate, Nottingham NG1 1LE (GB).</p>		<p>(81) Designated States: AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, JP, KE, KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MN, MW, MX, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TT, UA, UG, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ, UG).  <b>Published</b> <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>
<p>(54) Title: YEAST STRAINS AND MODIFIED ALBUMINS</p> <p>(57) Abstract</p> <p>Albumin, for example human albumin, is expressed and secreted in yeast which has been mutated to lack the yeast aspartyl protease 3 (Yap3p) or its equivalent, thereby reducing the production of a 45kD albumin fragment. A further reduction is achieved by additionally deleting the Kex2p function. Alternatively, a modified albumin is prepared which is not susceptible to Yap3p cleavage, for example human albumin which is R410A, K413Q and K414Q.</p> <div data-bbox="396 714 795 1155"> </div>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

## YEAST STRAINS AND MODIFIED ALBUMINS

### Field of the invention


- 5 The present invention relates to the production of recombinant human albumin (rHA) by yeast species.

### Background and prior art

- 10 Human serum albumin (HSA) is a protein of 585 amino acids that is responsible for a significant proportion of the osmotic pressure of serum, and also functions as a carrier of endogenous and exogenous ligands. It is used clinically in the treatment of patients with severe burns, shock, or blood loss, and at present is produced commercially by extraction from human blood. The  
15 production of recombinant human albumin (rHA) in microorganisms has been disclosed in EP 330 451 and EP 361 991.

- In recent years yeast species have been widely used as a host organisms for the production of heterologous proteins (reviewed by Romanos *et al*, 1992),  
20 including rHA (Sleep *et al*, 1990, 1991; Fleer *et al*, 1991). Yeasts are readily amenable to genetic manipulation, can be grown to high cell density on simple media, and as eukaryotes are suitable for production of secreted as well as cytosolic proteins.
- 25 When *S. cerevisiae* is utilised to produce rHA, the major secreted protein is mature 67kDa albumin. However, a 45kDa N-terminal fragment of rHA is also observed (Sleep *et al*, 1990). A similar fragment is obtained when rHA is expressed in *Kluyveromyces* sp. (Fleer *et al*, 1991) and *Pichia pastoris* (EP 510 693). The fragment has the same N-terminal amino acid sequence as  
30 mature rHA, but the carboxy terminus is heterogeneous and occurs between

Phe<sup>403</sup> and Val<sup>409</sup> with the most common termini being Leu<sup>407</sup> and Val<sup>409</sup> (Geisow *et al*, 1991), as shown below.



The amount of fragment produced, as a percentage of total rHA secreted, varies with both the strain and the secretion leader sequence utilised, but is never reduced to zero (Sleep *et al*, 1990). We have also found that the amount of fragment produced in high cell density fermentation (75-100g/L cell dry weight) is approximately five times higher than in shake flask cultures.

The 45kDa albumin fragment is not observed in serum-derived human serum albumin (HSA), and its presence as non-nature-identical material in the recombinant product is undesirable. The problem addressed by the present invention is to reduce the amount of the 45kDa fragment in the product. The simplest and most obvious approach would have been to have purified it away from the full length albumin, as proposed by Gist-brocades in EP 524 681 (see especially page 4, lines 17-22). However, we have chosen a different approach, namely to try to avoid its production in the first place.

Sleep *et al* (1990) postulated that rHA fragment is produced within the cell and is not the result of extra-cellular proteolysis. These authors codon-optimised the HSA cDNA from Glu<sup>382</sup> to Ser<sup>419</sup> but this had no effect on production of rHA fragment. They noted that a potential Kex2p processing site in the rHA amino acid sequence, Lys<sup>413</sup>Lys<sup>414</sup>, is in close proximity to the heterogeneous carboxy terminus of the fragment, but neither use of a *kex2* host strain (ie a strain harbouring a mutation in the *KEX2* gene such that it does not produce the Kex2p protease), nor removal of the potential cleavage site by site-directed

mutagenesis of the codon for Lys<sup>414</sup>, resulted in reduction in the amount of the fragment.

There is a vast array of yeast proteases which could, in principle, be degrading a desired protein product, including (in *S. cerevisiae*) yscA, yscB, yscY, yscS, other vacuolar proteinases, yscD, yscE, yscF (equivalent to kex2p), ysc $\alpha$ , yscIV, yscG, yscH, yscJ, yscE and kex1.

Bourbonnais *et al* (1991) described an *S. cerevisiae* endoprotease activity specific for monobasic sites, an example of which (Arg<sup>410</sup>) exists in this region of albumin. This activity was later found to be attributable to yeast aspartyl protease 3 (Yap3) (Bourbonnais *et al*, 1993), an enzyme which was originally described by Egel-Mitani *et al* (1990) as an endoprotease similar to Kex2p in specificity, in that it cleaved at paired basic residues. Further work suggested that Yap3p is able to cleave monobasic sites and between, and C-terminal to, pairs of basic residues, but that cleavage at both types of sites is dependent on the sequence context (Azaryan *et al*, 1993; Cawley *et al*, 1993).

As already discussed, the region of the C-terminus of rHA fragment contains both a monobasic (Arg<sup>410</sup>) and a dibasic site (Lys<sup>413</sup>Lys<sup>414</sup>). However, even though a Kex2p-like proteolytic activity is present in human cells and is responsible for cleavage of the pro sequence of HSA C-terminal to a pair of arginine residues, the fragment discussed above is not known to be produced in humans. This indicates that the basic residues Arg<sup>410</sup>, Lys<sup>413</sup> and Lys<sup>414</sup> are not recognised by this Kex2p-like protease, in turn suggesting that this region of the molecule may not be accessible to proteases in the secretory pathway. Thus, the Yap3p protease could not have been predicted to be responsible for the production of the 45kDa fragment. In addition, Egel-Mitani *et al* (1990 *Yeast* 6, 127-137) had shown Yap3p to be similar to Kex2p in cleaving the MF $\alpha$  propheromone. Since removal of the Kex2p function alone does not

reduce the amount of the fragment produced, there was no reason to suppose that removal of the Yap3p function would be beneficial. Indeed, Bourbonnais *et al* (1993) showed that *yap3* strains had a decreased ability to process pro-somatostatin, and therefore taught away from using *yap3* strains in the production of heterologous proteins.

### Summary of the invention

The solution to the problem identified above is, in accordance with the invention, to avoid or at least reduce production of the fragment in the initial fermentation, rather than to remove it during purification of the albumin. We have now found that, out of the 20 or more yeast proteases which are so far known to exist, it is in fact the Yap3p protease which is largely responsible for the 45kD fragment of rHA produced in yeast. The present invention provides a method for substantially reducing the amount of a 45kDa fragment produced when rHA is secreted from yeast species. The reduction in the amount of fragment both improves recovery of rHA during the purification process, and provides a higher quality of final product. A further, and completely unexpected, benefit of using *yap3* strains of yeast is that they can produce 30-50% more rHA than strains having the Yap3p function. This benefit cannot be accounted for merely by the reduction of rHA fragment from ~ 15% to 3-5%.

Thus, one aspect of the present invention provides a process for preparing albumin by secretion from a yeast genetically modified to produce and secrete the albumin, comprising culturing the yeast in a culture medium such that albumin is secreted into the medium, characterised in that the yeast cells have a reduced level of yeast aspartyl protease 3 proteolytic activity.

Preferably, the said proteolytic activity is an endoprotease activity specific for monobasic sites and for paired basic amino acids in a polypeptide.

Suitably, the yeast is *S. cerevisiae* which lacks a functional *YAP3* gene. However, the invention is not limited to the use of *S. cerevisiae*, since the problem of 45 kDa fragment production is found also in other yeast genera, for example *Pichia* and *Kluyveromyces*, which shows that they have equivalent proteases (ie Yap3p proteolytic activity); see Clerc *et al* (1994), page 253. We have confirmed this by hybridisation analysis to locate homologues of Yap3p in non-*Saccharomyces* genera. A gene is regarded as a homologue, in general, if the sequence of the translation product has greater than 50% sequence identity to Yap3p. In non-*Saccharomyces* genera, the Yap3p-like protease and its gene may be named differently, but this does not of course alter their essential nature.

The level of fragment can be reduced still further if, as well as substantially eliminating the Yap3p proteolytic activity, the Kex2p function is also substantially eliminated even though, as mentioned above, elimination of the Kex2p function alone does not affect the level of fragment. As in the case of Yap3p, the Kex2p function is not restricted to *Saccharomyces*; see Gellissen *et al* (1992), especially the sentence bridging pages 415 and 416, showing that *Pichia* has a Kex2p function. The genes encoding the Kex2p equivalent activity in *Kluyveromyces lactis* and *Yarrowia lipolytica* have been cloned (Tanguy-Rougeau *et al*, 1988; Enderlin & Ogrydziak, 1994).

A suitable means of eliminating the activity of a protease is to disrupt the host gene encoding the protease, thereby generating a non-reverting strain missing all or part of the gene for the protease (Rothstein, 1983). Alternatively, the activity can be reduced or eliminated by classical mutagenesis procedures or by the introduction of specific point mutations by the process of transplacement (Winston *et al*, 1983). Preferably, the activity of the enzyme is reduced to at most 50% of the wild-type level, more preferably no more than 25%, 10% or 5%, and most preferably is undetectable. The level of Yap3p proteolytic

activity may be measured by determining the production of the 45 kDa fragment, or by the  $^{125}\text{I}$ - $\beta_2$ -lipoprotein assay of Azaryan *et al* (1993), also used by Cawley *et al* (1993). Kex2p proteolytic activity may similarly be measured by known assays, for example as set out in Fuller *et al* (1989).

5

The albumin may be a human albumin, or a variant thereof, or albumin from any other animal.

By "variants" we include insertions, deletions and substitutions, either conservative or non-conservative, where such changes do not substantially alter the oncotic, useful ligand-binding or non-immunogenic properties of albumin. In particular, we include naturally-occurring polymorphic variants of human albumin; fragments of human albumin which include the region cleaved by Yap3p, for example those fragments disclosed in EP 322 094 (namely HSA (1-n), where n is 369 to 419) which are sufficiently long to include the Yap3p-cleaved region (ie where n is 403 to 419); and fusions of albumin (or Yap3p-cleavable portions thereof) with other proteins, for example the kind disclosed in WO 90/13653.

20 By "conservative substitutions" is intended swaps within groups such as Gly, Ala; Val, Ile, Leu; Asp, Glu; Asn, Gln; Ser, Thr; Lys, Arg; and Phe, Tyr.

Such variants may be made using the methods of protein engineering and site-directed mutagenesis as described below.

25

A second aspect of the invention provides a modified albumin having at least 90% sequence identity to a naturally-occurring albumin, which naturally-occurring albumin is susceptible to cleavage with the *S. cerevisiae* yeast aspartyl protease 3 (Yap3p) when expressed in yeast, characterised in that the modified albumin is not susceptible to such cleavage.



Preferably, the modified albumin lacks a monobasic amino acid present in the naturally-occurring albumin protein. Suitably, the said monobasic amino acid is arginine. Conveniently, the modified albumin additionally lacks a pair of basic amino acids present in the naturally-occurring albumin, especially any of

5 Lys, Lys; Lys, Arg; Arg, Lys; or Arg, Arg. Thus, in one particular embodiment, the naturally-occurring albumin is human albumin and the modified protein lacks Arg<sup>410</sup> and, optionally, one or both Lys<sup>413</sup>Lys<sup>414</sup> lysines. For example, the modified albumin may be human albumin having the amino acid changes R410A, K413Q, K414Q. Equivalent modifications in bovine

10 serum albumin include replacing the Arg<sup>408</sup> and/or one or both of Arg<sup>411</sup>Lys<sup>412</sup>. The person skilled in the art will be able to identify monobasic sites and pairs of basic residues in other albumins without difficulty.

The numbering of the residues corresponds to the sequence of normal mature

15 human albumin. If the albumin is a variant (for example a polymorphic form) having a net deletion or addition of residues N-terminal to the position identified, then the numbering refers to the residues of the variant albumin which are aligned with the numbered positions of normal albumin when the two sequences are so aligned as to maximise the apparent homology.

20

A third aspect of the invention provides a polynucleotide encoding such a modified albumin.

The DNA is expressed in a suitable yeast (either the DNA being for a modified

25 albumin, or the yeast lacking the Yap3p function) to produce an albumin. Thus, the DNA encoding the albumin may be used in accordance with known techniques, appropriately modified in view of the teachings contained herein, to construct an expression vector, which is then used to transform an appropriate yeast cell for the expression and production of the albumin.

30

The DNA encoding the albumin may be joined to a wide variety of other DNA sequences for introduction into an appropriate host. The companion DNA will depend upon the nature of the host, the manner of the introduction of the DNA into the host, and whether episomal maintenance or integration is desired.

5

Generally, the DNA is inserted into an expression vector, such as a plasmid, in proper orientation and correct reading frame for expression. The vector is then introduced into the host through standard techniques and, generally, it will be necessary to select for transformed host cells.

10

Host cells that have been transformed by the recombinant DNA of the invention are then cultured for a sufficient time and under appropriate conditions known to those skilled in the art in view of the teachings disclosed herein to permit the expression and secretion of the albumin, which can then be recovered, as is known.

15

Useful yeast plasmid vectors are pRS403-406 and pRS413-416 and are generally available from Stratagene Cloning Systems, La Jolla, CA 92037, USA. Plasmids pRS403, pRS404, pRS405 and pRS406 are Yeast Integrating plasmids (YIps) and incorporate the yeast selectable markers *HIS3*, *TRP1*, *LEU2* and *URA3*. Plasmids pRS413-416 are Yeast Centromere plasmids (YCps). Other yeast expression plasmids are disclosed in EP-A-258 067, EP-A-286 424 and EP-A-424 117.

20

The polynucleotide coding sequences encoding the modified albumin of the invention may have additional differences to those required to produce the modified albumin. For example, different codons can be substituted which code for the same amino acid(s) as the original codons. Alternatively, the substitute codons may code for a different amino acid that will not affect the activity or immunogenicity of the albumin or which may improve its activity

30

or immunogenicity, as well as reducing its susceptibility to a Yap3p protease activity. For example, site-directed mutagenesis or other techniques can be employed to create single or multiple mutations, such as replacements, insertions, deletions, and transpositions, as described in Botstein and Shortle (1985). Since such modified coding sequences can be obtained by the application of known techniques to the teachings contained herein, such modified coding sequences are within the scope of the claimed invention.

Exemplary genera of yeast contemplated to be useful in the practice of the present invention are *Pichia*, *Saccharomyces*, *Kluyveromyces*, *Candida*, *Torulopsis*, *Hansenula* (now reclassified as *Pichia*), *Histoplasma*, *Schizosaccharomyces*, *Citeromyces*, *Pachysolen*, *Debaromyces*, *Metschnikowia*, *Rhodospordium*, *Leucosporidium*, *Botryosascus*, *Sporidiobolus*, *Endomycopsis*, and the like. Preferred genera are those selected from the group consisting of *Pichia*, *Saccharomyces*, *Kluyveromyces*, *Yarrowia* and *Hansenula*. Examples of *Saccharomyces* sp. are *S. cerevisiae*, *S. italicus* and *S. rouxii*. Examples of *Kluyveromyces* sp. are *K. fragilis* and *K. lactis*. Examples of *Hansenula* (*Pichia*) sp. are *H. polymorpha* (now *Pichia angusta*), *H. anomala* (now *P. anomala*) and *P. pastoris*. *Y. lipolytica* is an example of a suitable *Yarrowia* species.

Methods for the transformation of *S. cerevisiae* are taught generally in EP 251 744, EP 258 067 and WO 90/01063, all of which are incorporated herein by reference. Suitable promoters for *S. cerevisiae* include those associated with the *PGK1* gene, *GAL1* or *GAL10* genes, *CYC1*, *PHO5*, *TRP1*, *ADH1*, *ADH2*, the genes for glyceraldehyde-3-phosphate dehydrogenase, hexokinase, pyruvate decarboxylase, phosphofructokinase, triose phosphate isomerase, phosphoglucose isomerase, glucokinase,  $\alpha$ -mating factor pheromone, a-mating factor pheromone, the *PRB1* promoter, the *GPD1* promoter, and hybrid promoters involving hybrids of parts of 5' regulatory regions with parts of 5'

regulatory regions of other promoters or with upstream activation sites (eg the promoter of EP-A-258 067).

- Convenient regulatable promoters for use in *Schizosaccharomyces pombe* are
- 5 the thiamine-repressible promoter from the *nmt* gene as described by Maundrell (1990) and the glucose-repressible *fbp1* gene promoter as described by Hoffman & Winston (1990).

- Methods of transforming *Pichia* for expression of foreign genes are taught in,
- 10 for example, Cregg *et al* (1993), and various Phillips patents (eg US 4 857 467, incorporated herein by reference), and *Pichia* expression kits are commercially available from Invitrogen BV, Leek, Netherlands, and Invitrogen Corp., San Diego, California. Suitable promoters include *AOX1* and *AOX2*.

- 15 The Gellissen *et al* (1992) paper mentioned above and Gleeson *et al* (1986) *J. Gen. Microbiol.* **132**, 3459-3465 include information on *Hansenula* vectors and transformation, suitable promoters being *MOX1* and *FMD1*; whilst EP 361 991, Fleer *et al* (1991) and other publications from Rhône-Poulenc Rorer teach how to express foreign proteins in *Kluyveromyces* spp., a suitable promoter being
- 20 *PGK1*.

- The transcription termination signal is preferably the 3' flanking sequence of a eukaryotic gene which contains proper signals for transcription termination and polyadenylation. Suitable 3' flanking sequences may, for example, be
- 25 those of the gene naturally linked to the expression control sequence used, ie may correspond to the promoter. Alternatively, they may be different in which case the termination signal of the *S. cerevisiae ADH1* gene is preferred.

- The albumin is initially expressed with a secretion leader sequence, which may
- 30 be any leader effective in the yeast chosen. Leaders useful in *S. cerevisiae*

include that from the mating factor  $\alpha$  polypeptide (MF $\alpha$ -1) and the hybrid leaders of EP-A-387 319. Such leaders (or signals) are cleaved by the yeast before the mature albumin is released into the surrounding medium. When the yeast strain lacks Kex2p activity (or equivalent) as well as being *yap3*, it may be advantageous to choose a secretion leader which need not be cleaved from the albumin by Kex2p. Such leaders include those of *S. cerevisiae* invertase (*SUC2*) disclosed in JP 62-096086 (granted as 91/036516), acid phosphatase (*PHO5*), the pre-sequence of MF $\alpha$ -1,  $\beta$ -glucanase (*BGL2*) and killer toxin; *S. diastaticus* glucoamylase II; *S. carlsbergensis*  $\alpha$ -galactosidase (*MEL1*); *K. lactis* killer toxin; and *Candida* glucoamylase.

Various non-limiting embodiments of the invention will now be described by way of example and with reference to the accompanying drawings in which:

Figure 1 is a general scheme for the construction of mutated rHA expression plasmids, in which HA is a human albumin coding sequence, L is a sequence encoding a secretion leader, P is the PRB1 promoter, T is the ADH1 terminator, amp is an ampicillin resistance gene and LEU2 is the leucine selectable marker;

Figure 2 is a drawing representing a Western blot analysis of mutant rHA secreted by *S. cerevisiae*, in which Track A represents the culture supernatant from DB1 cir<sup>-</sup> pAYE316 (normal rHA), Track B represents the culture supernatant from DB1 cir<sup>+</sup> pAYE464 (alteration 1), and Track C represents the culture supernatant from DB1 cir<sup>+</sup> pAYE468 (alteration 3);

Figure 3 is a scheme of the construction of pAYE515;

Figure 4 is a comparison of rHA fragment production by wild-type and protease-disrupted strains, presented as a drawing of an anti-HSA Western blot

of culture supernatant from shake flask cultures separated by non-reducing 10% SDS/PAGE, in which Track A corresponds to DB1 cir<sup>o</sup> pAYE316, Track B corresponds to DXY10 cir<sup>o</sup> pAYE316 (*yap3* strain), and Track C corresponds to ABB50 cir<sup>o</sup> pAYE316 (*yap3*, *kex2* strain);

5

Figure 5 is similar to Figure 4 but shows Coomassie Brilliant Blue stained 12.5% SDS Phastgel (Pharmacia) of culture supernatants from fed batch fermentations, namely Track D for the HSA standard, Track E for DB1 cir<sup>o</sup> pAYE316, Track F for DB1  $\Delta kex2$  cir<sup>o</sup> pAYE522, and Track G for DXY10 cir<sup>o</sup> pAYE522; and

10

Figure 6 is a scheme for the construction of pAYE519.

#### Detailed description of the invention

15

All standard recombinant DNA procedures are as described in Sambrook *et al* (1989) unless otherwise stated. The DNA sequences encoding HSA are derived from the cDNA disclosed in EP 201 239.

#### 20 **Example 1: Modification of the HSA cDNA.**

In order to investigate the role of endoproteases in the generation of rHA fragment, the HSA cDNA (SEQ1 (which includes a sequence encoding the artificial secretion leader sequence of WO 90/01063)) was modified by site-directed mutagenesis. Three separate changes were made to the HSA sequence (SEQ2). The first, using the mutagenic primer FOG1, changed the Arg<sup>410</sup> codon only, replacing it with an Ala codon, leaving intact the dibasic site, Lys<sup>413</sup>Lys<sup>414</sup>. The second change, using primer FOG2, changed the residues 407-409, including the C-terminal residues of fragment, from LeuLeuVal to AlaValAla. The third change, using the primer FOG3, altered residues 410-

25

30

414 from ArgTyrThrLysLys (SEQ3) to AlaTyrThrGlnGln (SEQ4). The oligonucleotides encoded not only the amino acid changes, but also conservative base changes that create either a *PvuII* or an *SpeI* restriction site in the mutants to facilitate detection of the changed sequences.

5

Single-stranded DNA of an M13mp19 clone, mp19.7 (EP 201 239; Figure 2), containing the HSA cDNA was used as the template for the mutagenesis reactions using the *In Vitro* Mutagenesis System, Version 2 (Amersham International plc) according to the manufacturer's instructions. Individual  
10 plaques were selected and sequenced to confirm the presence of the mutations. Double stranded RF DNA was then made from clones with the expected changes and the DNA bearing the mutation was excised on an *XbaI/SacI* fragment (Figure 1). This was used to replace the corresponding wild-type fragment of pAYE309 (EP 431 880; Figure 2). The presence of the mutated  
15 *XbaI/SacI* fragment within the plasmid was checked by digesting with *PvuII* or *SpeI* as appropriate. These *HindIII* fragments were excised and inserted into the expression vector pAYE219 (Figure 1) to generate the plasmids pAYE464 (alteration 1, R410A), pAYE470 (alteration 2, L407A, L408V, V409A) and pAYE468 (alteration 3, R410A, K413Q, K414Q). These expression plasmids  
20 comprise the *S. cerevisiae PRB1* promoter (WO 91/02057) driving expression of the HSA/MF $\alpha$ 1 leader sequence (WO 90/01063) fused in-frame with the mutated HA coding sequence which is followed by the *ADHI* transcription terminator. The plasmids also contain part of the 2 $\mu$ m plasmid to provide replication functions and the *LEU2* gene for selection of transformants.

25

pAYE464, pAYE470 and pAYE468 were introduced into *S. cerevisiae* DB1 *cir*<sup>+</sup> (*a. leu2*; Sleep *et al*, 1990) by transformation and individual transformants were grown for 3 days at 30°C in 10ml YEPS (1% w/v yeast extract, 2% w/v peptone, 2% w/v sucrose) and then the supernatants were examined by anti-HSA  
30 Western blot for the presence of the rHA fragment. The Western blots clearly

showed that fragment was still produced by the strains harbouring pAYE464, although the level was reduced slightly compared to the control expressing wild-type rHA. The mutations in the plasmid pAYE470 appeared to have no effect on the generation of fragment. However, DB1 cir<sup>+</sup> pAYE468 showed a novel pattern of HSA-related bands, with little or no fragment.

One example of each of DB1 cir<sup>+</sup> pAYE464 and DB1 cir<sup>+</sup> pAYE468 were grown to high cell density by fed batch culture in minimal medium in a fermenter (Collins, 1990). Briefly, a fermenter of 10L working volume was filled to 5L with an initial batch medium containing 50 mL/L of a concentrated salts mixture (Table 1), 10 mL/L of a trace elements solution (Table 2), 50 mL/L of a vitamins mixture (Table 3) and 20 g/L sucrose. An equal volume of feed medium containing 100 mL/L of the salts mixture, 20 mL/L of the trace elements mixture, 100 mL/L of vitamins solution and 500 g/L sucrose was held in a separate reservoir connected to the fermenter by a metering pump. The pH was maintained at  $5.7 \pm 0.2$  by the automatic addition of ammonium hydroxide or sulphuric acid, and the temperature was maintained at 30°C. The stirrer speed was adjusted to give a dissolved oxygen tension of >20% air saturation at 1 v/v/min air flow rate.

**Table 1. Salts Mixture**

Chemical	Concentration (g/L)
KH <sub>2</sub> PO <sub>4</sub>	114.0
MgSO <sub>4</sub>	12.0
CaCl <sub>2</sub> .6H <sub>2</sub> O	3.0
Na <sub>2</sub> EDTA	2.0



Table 2. Trace Elements Solution

Chemical	Concentration (g/L)
ZnSO <sub>4</sub> ·7H <sub>2</sub> O	3.0
FeSO <sub>4</sub> ·7H <sub>2</sub> O	10.0
MnSO <sub>4</sub> ·4H <sub>2</sub> O	3.2
CuSO <sub>4</sub> ·5H <sub>2</sub> O	0.079
H <sub>3</sub> BO <sub>3</sub>	1.5
KI	0.2
Na <sub>2</sub> MoO <sub>4</sub> ·2H <sub>2</sub> O	0.5
CoCl <sub>2</sub> ·6H <sub>2</sub> O	0.56
H <sub>3</sub> PO <sub>4</sub>	75mL/L

Table 3. Vitamins Solution

Chemical	Concentration (g/L)
Ca pantothenate	1.6
Nicotinic acid	1.2
m inositol	12.8
Thiamine HCl	0.32
Pyridoxine HCl	0.8
Biotin	0.008

The fermenter was inoculated with 100 mL of an overnight culture of *S. cerevisiae* grown in buffered minimal medium (Yeast nitrogen base [without amino acids, without ammonium sulphate, Difco] 1.7 g/L, (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub> 5 g/L, citric acid monohydrate 6.09 g/L, Na<sub>2</sub>HPO<sub>4</sub> 20.16 g/L, sucrose 20 g/L, pH6.5). The initial batch fermentation proceeded until the carbon source had been consumed, at which point the metering pump was switched on and the addition of feed was computer controlled (the micro MFCS system, B. Braun,

Melsungen, Germany) using an algorithm based on that developed by Wang *et al* (1979). A mass spectrometer was used in conjunction with the computer control system to monitor the off gases from the fermentation and to control the addition of feed to maintain a set growth rate (eg 0.1 h<sup>-1</sup>). Maximum conversion of carbon substrate into biomass is achieved by maintaining the respiratory coefficient below 1.2 (Collins, 1990) and, by this means, cell densities of approximately 100 g/L cell dry weight can be achieved. The culture supernatants were compared with those of a wild-type rHA producer by Coomassie-stained SDS/PAGE and by Western blot. These indicated (Figure 2) that, whilst elimination of the monobasic Arg<sup>410</sup> (pAYE464) did reduce the level of the fragment by a useful amount, removal of both potential protease sites (pAYE468) almost abolished the 45kDa fragment.

The above data suggested that the generation of rHA fragment might be due to endoproteolytic attack, though the absence of an effect of removal of the potential Kex2p site Lys<sup>413</sup>Lys<sup>414</sup> (Sleep *et al*, 1990, and confirmed by other studies not noted here) unless combined with elimination of Arg<sup>410</sup>, had suggested a complex etiology. The reduction in the amount of fragment with the mutated rHA could in principle be due to an effect of the changes on the kinetics of folding of the molecule and not due to the removal of protease cleavage sites.

#### Example 2: Disruption of the YAP3 gene.

The YAP3 gene encoding yeast aspartyl protease 3 was mutated by the process of gene disruption (Rothstein 1983) which effectively deleted part of the YAP3 coding sequence, thereby preventing the production of active Yap3p.

Four oligonucleotides suitable for PCR amplification of the 5' and 3' ends of the YAP3 gene (Egel-Mitani *et al*, 1990) were synthesised using an Applied

Biosystems 380B Oligonucleotide Synthesiser. To assist the reader, we include as SEQ15 the sequence of the *YAP3* gene, of which 541-2250 is the coding sequence.

## 5 5' end

YAP3A: 5'-CGTCAGACCTTGCATGCAGCCAAGACACCCTCACATAGC-3' (SEQ5)

YAP3B: 5'-CCGTTACGTTCTGTGGTGGCATGCCCACTTCCAAGTCCACCG-3' (SEQ6)

10

## 3' end

YAP3C: 5'-GCGTCTCATAGTGGAAAAGCTTCTAAATACGACAACTTCCCC-3' (SEQ7)

15 YAP3D: 5'-CCCAAAATGGTACCTGTGTCACTCGTTGGGATAATACC-3' (SEQ8)

PCR reactions were carried out to amplify individually the 5' and 3' ends of the *YAP3* gene from *S. cerevisiae* genomic DNA (Clontech Laboratories, Inc). Conditions were as follows: 2.5µg/ml genomic DNA, 5µg/ml of each primer, 20 denature at 94°C 61 seconds, anneal at 37°C 121 secs, extend at 72°C 181 secs for 40 cycles, followed by a 4°C soak, using a Perkin-Elmer-Cetus Thermal Cycler and a Perkin-Elmer-Cetus PCR kit according to the manufacturer's recommendations. Products were analysed by gel electrophoresis and were found to be of the expected size. The 5' fragment 25 was digested with *SphI* and cloned into the *SphI* site of pUC19HX (pUC19 lacking a *HindIII* site) to give pAYE511 (Figure 3), in which the orientation is such that *YAP3* would be transcribed towards the *KpnI* site of the pUC19HX polylinker. The 3' *YAP3* fragment was digested with *HindIII* and *Asp718* (an isoschizomer of *KpnI*) and ligated into pUC19 digested with *HindIII*/*Asp718* to 30 give pAYE512. Plasmid DNA sequencing was carried out on the inserts to confirm that the desired sequences had been cloned. The *HindIII*/*Asp718* fragment of pAYE512 was then subcloned into the corresponding sites of pAYE511 to give pAYE513 (Fig 3), in which the 5' and 3' regions of *YAP3*

are correctly orientated with a unique *Hind*III site between them. The *URA3* gene was isolated from YEp24 (Botstein *et al*, 1979) as a *Hind*III fragment and then inserted into this site to give pAYE515 (Fig 3), with *URA3* flanked by the 5' and 3' regions of *YAP3*, and transcribed in the opposite direction to *YAP3*.

5

A *ura3* derivative of strain DB1 cir<sup>o</sup> pAYE316 (Sleep *et al*, 1991) was obtained by random chemical mutagenesis and selection for resistance to 5-fluoro-orotic acid (Boeke *et al*, 1987). The strain was grown overnight in 100 mL buffered minimal medium and the cells were collected by centrifugation and then washed once with sterile water. The cells were then resuspended in 10 mL sterile water and 2 mL aliquots were placed in separate 15 mL Falcon tubes. A 5 mg/mL solution of N-methyl-N'-nitro-N-nitrosoguanidine (NTG) was then added to the tubes as follows: 0  $\mu$ L, 20  $\mu$ L, 40  $\mu$ L, 80  $\mu$ L or 160  $\mu$ L. The cells were then incubated at 30°C for 30 min and then centrifuged and washed three times with sterile water. Finally, the cells were resuspended in 1 mL YEP (1% w/v yeast extract, 2% w/v Bacto peptone) and stored at 4°C. The percentage of cells that survived the mutagenic treatment was determined by spreading dilutions of the samples on YEP plates containing 2% w/v sucrose and incubating at 30°C for 3 days. Cells from the treatment which gave approximately 50% survival were grown on YEP plates containing 2% w/v sucrose and then replica-plated onto YNB minimal medium containing 2% w/v sucrose and supplemented with 5-fluoro-orotic acid (1 mg/mL) and uracil (50  $\mu$ g/mL). Colonies able to grow on this medium were purified, tested to verify that they were unable to grow in the absence of uracil supplementation and that this defect could be corrected by introduction of the *URA3* gene by transformation. One such strain, DBU3 cir<sup>o</sup> pAYE316, was transformed with the *Sph*I/*Asp*718 *YAP3-URA3-YAP3* fragment of pAYE515 with selection for Ura<sup>+</sup> colonies. A Southern blot of digested genomic DNA of a number of transformants was probed with the 5' and 3' ends of the *YAP3* gene and confirmed the disruption of the *YAP3* gene. An anti-HSA Western blot of

30

YEPS shake-flask supernatants of two transformants indicated that disruption of *YAP3* markedly reduced rHA fragment levels.

- One *yap3* derivative of DBU3 cir° pAYE316, designated DXY10 cir° pAYE316, was grown several times by fed-batch fermentation in minimal medium to high cell dry weight. When supernatants were examined by Coomassie-stained PAGE and anti-HSA Western blot (Figs 4 and 5), the reduction in the level of rHA 45kDa fragment was clearly apparent; estimates of the amount of the degradation product vary from  $1/3$  to  $1/5$  of the levels seen with the *YAP3* parent. The amount of rHA produced was not adversely affected by the *yap3* mutation, indeed DXY10 cir° pAYE316 was found to produce 30-50% more rHA than the *YAP3* equivalent, DB1 cir° pAYE316. Despite the fact that cleavage of the leader sequence from the HA sequence is C-terminal to a pair of basic residues, the rHA was found to have the correct N-terminus.

- The fermentation broth was centrifuged to remove the cells and then subject to affinity chromatographic purification as follows. The culture supernatant was passed through a Cibacron Blue F3GA Sepharose column (Pharmacia) which was then washed with 0.1M phosphate glycine buffer, pH8.0. The rHA was then eluted from the column with 2M NaCl, 0.1M phosphate glycine, pH8.0, at which point it was >95% pure. It may be purified further by techniques known in the art.

- The albumin may alternatively be purified from the culture medium by any of the variety of known techniques for purifying albumin from serum or fermentation culture medium, for example those disclosed in WO 92/04367, Maurel *et al* (1989), Curling (1980) and EP 524 681.

**Example 3: Disruption of the *KEX2* gene in a *yap3* strain.**

To construct a strain lacking both Yap3p and Kex2p activity, a *lys2* derivative of yeast strain DXY10 cir<sup>o</sup> (pAYE316) was obtained by random chemical mutagenesis and selection for resistance to  $\alpha$ -amino adipate (Barnes and Thorner, 1985). Cells were mutagenised as in Example 2 and then plated on YNB minimal medium containing 2% w/v sucrose and supplemented with 2 mg/mL DL- $\alpha$ -amino adipate as the sole nitrogen source and 30  $\mu$ g/mL lysine. Colonies able to grow on this medium were purified and tested to verify that they were unable to grow in the absence of lysine supplementation and that this defect could be corrected by the introduction of the *LYS2* gene by transformation. This strain was then mutated by the process of gene disruption which effectively disrupted part of the *KEX2* coding sequence, thereby preventing production of active Kex2p. To assist the reader, the sequence of the *KEX2* gene is reproduced herein as SEQ14, of which 1329-3773 is the coding sequence.

Four oligonucleotides suitable for PCR amplification of the 5' and 3' ends of the *KEX2* gene (Fuller *et al*, 1989) were synthesised using an Applied Biosystems 380B Oligonucleotide Synthesiser.

**5' end**

KEX2A: 5'-CCATCTGGATCCAATGGTGCTTTGGCCAAATAAATAGTTTCAGC-3' (SEQ9)  
KEX2B: 5'-GCTTCTTTTACCGGTAACAAGCTTGAGTCCATTGG-3' (SEQ10)

**3' end**

KEX2C: 5'-GGTAAGGTTTACTCGACCTATTTTTTGTTTTGTCTGC-3' (SEQ11)  
KEX2D: 5'-GGAAACGTATGAATTCGATATCATTGATACAGACTCTGAGTACG-3' (SEQ12)

PCR reactions were carried out to amplify individually the 5' and 3' ends of the *KEX2* gene from *S. cerevisiae* genomic DNA (Clontech Laboratories Inc). Conditions were as follows: 2.5 µg/ml genomic DNA, 5 µg/ml of each primer, denature 94°C 61s, anneal 37°C 121s, extend 72°C 181s for 40 cycles, followed by a 4°C soak, using a Perkin-Elmer-Cetus Thermal Cycler and a Perkin-Elmer-Cetus PCR kit according to the manufacturer's recommendations. Products were analysed by gel electrophoresis and were found to be of the expected size (0.9 kb for the 5' product and 0.62 kb for the 3' product). The 5' product was digested with *Bam*HI and *Hind*III and the 3' product was digested with *Hind*III and *Sal*I and then the two fragments were together cloned into pUC19HX digested with *Bam*HI and *Sal*I. A 4.8 kb *Hind*III fragment comprising the *S. cerevisiae* *LYS2* gene (Barnes & Thorne, 1985) was then inserted into the resulting plasmid at *Hind*III (ie between the two *KEX2* fragments) to form pAYE519 (Fig 6).

The *lys2* derivative of DXY10 cir<sup>o</sup> (pAYE316), *lys2-16*, was transformed with the 6.0 kb *KEX2-LYS2-KEX2* fragment of pAYE519, selecting for Lys<sup>+</sup> colonies. A Southern blot of digested genomic DNA of a number of transformants was probed with the 5' and 3' ends of the *KEX2* gene and confirmed the disruption of the *KEX2* gene. An anti-HSA Western blot of YEPS shake-flask culture supernatants of these transformants indicated that disruption of *KEX2* in a *yap3* strain reduced the level of rHA fragment still further, despite the lack of an effect of disruption of *KEX2* alone in Example 4 below. Analysis of the rHA produced by one such strain, ABB50, indicated that the leader sequence was incorrectly processed, leading to an abnormal N-terminus.

The strain ABB50 (pAYE316) was cured of its plasmid (Sleep *et al*, 1991) and transformed with a similar plasmid, pAYE522, in which the hybrid leader sequence was replaced by the *S. cerevisiae* invertase (*SUC2*) leader sequence

such that the encoded leader and the junction with the HSA sequence were as follows:

	MLLQAFLFLLAGFAAKISA ↓ DAHKS	(SEQ13)
5	Invertase leader	HSA

In this construct, cleavage of the leader sequence from HSA does not rely upon activity of the Kex2 protease. The strain ABB50 (pAYE522) was found to produce rHA with a similarly very low level of rHA fragment, but in this instance the N-terminus corresponded to that of serum-derived HSA, ie there was efficient and precise removal of the leader sequence.

**Example 4: Disruption of the *KEX2* gene alone (Comparative Example).**

By a similar method to that disclosed in Example 3 the *KEX2* gene was disrupted in *S. cerevisiae*. This strain had the Yap3p proteolytic activity and was therefore not within the scope of the invention. When this strain was grown in fed batch fermentation the rHA produced contained similar amounts of fragment to that produced by strains with an intact *KEX2* gene. In addition, the overall level of rHA was reduced and the leader sequence was not correctly processed, leading to an abnormal N-terminus.

**Example 5: Identification of equivalent protease in *Pichia*.**

As noted above, non-*Saccharomyces* yeast similarly produce the undesirable fragment of rHA and therefore have the Yap3p proteolytic activity. We have confirmed this by performing Southern hybridisations of *Pichia angusta* DNA, using the *S. cerevisiae* *YAP3* gene as a probe. A specific DNA fragment was identified, showing that, not only is the Yap3p proteolytic activity present in *P. angusta*, but a specific homologue of the *YAP3* gene is present also.



The method of Southern hybridization used for detection of the *YAP3* homologue can be adapted to clone the gene sequence from a genomic DNA library of *Pichia* DNA using standard procedures (Sambrook *et al*, 1989). Disruption of the *YAP3* homologue in *Pichia sp.* can be achieved using similar  
5 techniques to those used above for *Saccharomyces* (Cregg and Madden, 1987).

## References (all incorporated by reference)

- Azaryan, A.V. *et al* (1993) *J. Biol. Chem.* **268**, 11968-11975.
- Barnes, D.A. and Thorner, J. (1985) *In* Gene Manipulations in Fungi (Bennett, J.W. and Lasure, L.L., eds), pp. 197-226, Academic Press.
- Boeke, J.D. *et al* (1987) *Methods Enzymol.* **154**, 164-175.
- Botstein, D. *et al* (1979) *Gene* **8**, 17-24.
- Botstein & Shortle (1985) *Science* **229**, 193-210.
- Bourbonnais, Y. *et al* (1991) *J. Biol. Chem.* **266**, 13203-13209.
- 10 Bourbonnais, Y. *et al* (1993) *EMBO J.* **12**, 285-294.
- Cawley, N.X. *et al* (1993) *FEBS Lett.* **332**, 273-276.
- Clerc *et al* (1994) *J. Chromat. B.* **662**, 245-259.
- Collins, S.H. (1990) *In* Protein Production by Biotechnology (Harris, T.J.R., ed) pp. 61-77, Elsevier Science Publishers, Barking, Essex.
- 15 Cregg, J.M. and Madden, K.R. (1987) *In* Biological Research on Industrial Yeasts, Vol. II, Stewart, G.G., Russell, I., Klein, R.D. and Hiebsch, R.R. (Eds) CRC Press, Boca Raton, FL.
- Cregg *et al* (1993) *Bio/Technology* **11**, 905-910.
- Curling (1980) "Albumin Purification by Ion Exchange Chromatography", in
- 20 "Methods of Plasma Protein Purification", Ed. Curling, J.M., Academic Press, London.
- Enderlin, C.S. & Ogrydziak, D.M. (1994) *Yeast* **10**, 67-79.
- Fleer, R. *et al* (1991) *Bio/Technology* **9**, 968-975.
- Fuller, R.S. *et al* (1989) *Proc. Natl. Acad. Sci. USA* **86**, 1434-1438.
- 25 Geisow, M.J. *et al* (1991) *In* Techniques in Protein Chemistry II, pp. 567-572, Academic Press, Inc.
- Gellissen *et al* (1992) *Tibtech* **10**, 413-417.
- Hoffmann & Winston (1990) *Genetics* **124**, 807-816.
- Maundrell (1990) *J. Biol. Chem.* **265**, 10857-10864.
- 30 Maurel *et al* (1989) "Biotechnology of Plasma Proteins", *Colloque INSERM*

- 175, 19-24.
- Romanos, M.A. (1992) *Yeast* **8**, 423-488.
- Rothstein, R.J. (1983) *Methods Enzymol.* **101**, 203-211.
- Sambrook, J. *et al* (1989) *Molecular Cloning: a Laboratory Manual*, 2nd  
5 edition. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
- Sleep, D. *et al* (1990) *Bio/Technology* **8**, 42-46.
- Sleep, D. *et al* (1991) *Bio/Technology* **9**, 183-187.
- Tanguy-Rougeau, C. *et al* (1988) *FEBS Lett.* **234**, 464-470
- Wang, H.Y. *et al* (1979) *Biotech. & Bioeng.* **21**, 975.
- 10 Winston, F. *et al* (1983) *Methods Enzymol.* **101**, 211-228.

## 26

## SEQUENCE LISTING

## (1) GENERAL INFORMATION:

## (i) APPLICANT:

- (A) NAME: Delta Biotechnology Limited
- (B) STREET: Castle Court, Castle Boulevard
- (C) CITY: Nottingham
- (D) STATE: Nottinghamshire
- (E) COUNTRY: United Kingdom
- (F) POSTAL CODE (ZIP): NG7 1FD

(ii) TITLE OF INVENTION: Yeast strains and modified albumins

(iii) NUMBER OF SEQUENCES: 15

## (iv) COMPUTER READABLE FORM:

- (A) MEDIUM TYPE: Floppy disk
- (B) COMPUTER: IBM PC compatible
- (C) OPERATING SYSTEM: PC-DOS/MS-DOS
- (D) SOFTWARE: Patentin Release #1.0, Version #1.25 (EPO)

## (2) INFORMATION FOR SEQ ID NO: 1:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1830 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA to mRNA

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

## (ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 73..1827

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

ATGAAGTGGG TAAGCTTTAT TTCCCTTCTT TTTCTCTTTA GCTCGGCTTA TTCAGGAGC	60
TTGGATAAAA GA GAT GCA CAC AAG AGT GAG GTT GCT CAT CGS TTT AAA	108
Asp Ala His Lys Ser Glu Val Ala His Arg Phe Lys	
1 5 10	
GAT TTG GGA GAA GAA AAT TTC AAA GCC TTG GTG TTG ATT GCC TTT GCT	156
Asp Leu Gly Glu Glu Asn Phe Lys Ala Leu Val Leu Ile Ala Phe Ala	
15 20 25	
CAG TAT CTT CAG CAG TGT CCA TTT GAA GAT CAT GTA AAA TTA GTG AAT	204
Gln Tyr Leu Gln Gln Cys Pro Phe Glu Asp His Val Lys Leu Val Asn	
30 35 40	
GAA GTA ACT GAA TTT GCA AAA ACA TGT GTT GCT GAT GAG TCA GCT GAA	252
Glu Val Thr Glu Phe Ala Lys Thr Cys Val Ala Asp Glu Ser Ala Glu	
45 50 55 60	

AAT Asn	TGT Cys	GAC Asp	AAA Lys	TCA Ser	CTT Leu	CAT His	ACC Thr	CTT Leu	TTT Phe	GGA Gly	GAC Asp	AAA Lys	TTA Leu	TGC Cys	ACA Thr	300
				65					70					75		
GTT Val	GCA Ala	ACT Thr	CTT Leu	CGT Arg	GAA Glu	ACC Thr	TAT Tyr	GGT Gly	GAA Glu	ATG Met	GCT Ala	GAC Asp	TGC Cys	TGT Cys	GCA Ala	348
			80					85					90			
AAA Lys	CAA Gln	GAA Pro	CCT Pro	GAG Glu	AGA Arg	AAT Asn	GAA Glu	TGC Cys	TTC Phe	TTG Leu	CAA Gln	CAC His	AAA Lys	GAT Asp	GAC Asp	396
			95				100					105				
AAC Asn	CCA Pro	AAC Asn	CTC Leu	CCC Pro	CGA Arg	TTG Leu	GTG Val	AGA Arg	CCA Pro	GAG Glu	GTT Val	GAT Asp	GTG Val	ATG Met	TGC Cys	444
			110			115					120					
ACT Thr	GCT Ala	TTT Phe	CAT His	GAC Asp	AAT Asn	GAA Glu	GAG Glu	ACA Thr	TTT Phe	TTG Leu	AAA Lys	AAA Lys	TAC Tyr	TTA Leu	TAT Tyr	492
			125		130				135					140		
GAA Glu	ATT Ile	GCC Ala	AGA Arg	AGA His	CAT Pro	CCT Tyr	TAC Phe	TTT Phe	TAT Tyr	GCC Ala	CCG Pro	GAA Glu	CTC Leu	CTT Leu	TTC Phe	540
				145				150						155		
TTT Phe	GCT Ala	AAA Lys	AGG Arg	TAT Tyr	AAA Lys	GCT Ala	GCT Ala	TTT Ala	ACA Thr	GAA Glu	TGT Cys	TGC Cys	CAA Gln	GCT Ala	GCT Ala	588
			160					165					170			
GAT Asp	AAA Lys	GCT Ala	GCC Ala	TGC Cys	CTG Leu	TTG Leu	CCA Pro	AAG Lys	CTC Leu	GAT Asp	GAA Glu	CTT Leu	CGG Arg	GAT Asp	GAA Glu	636
			175				180					185				
GGG Gly	AAG Ala	GCT Ser	TCG Ser	TCT Ala	GCC Ala	AAA Lys	CAG Gln	AGA Arg	CTC Leu	AAG Lys	TGT Cys	GCC Ala	AGT Ser	CTC Leu	CAA Gln	684
			190			195				200						
AAA Lys	TTT Phe	GGA Gly	GAA Glu	AGA Arg	GCT Ala	TTC Phe	AAA Lys	GCA Ala	TGG Trp	GCA Val	GTA Ala	GCT Arg	CGC Ala	CTG Leu	AGC Ser	732
			205		210				215					220		
CAG Gln	AGA Arg	TTT Phe	CCC Pro	AAA Lys	GCT Ala	GAG Glu	TTT Phe	GCA Ala	GAA Glu	GTT Val	TCC Ser	AAG Lys	TTA Leu	GTG Val	ACA Thr	780
				225				230					235			
GAT Asp	CTT Leu	ACC Thr	AAA Lys	GTG Val	CAC His	ACG Thr	GAA Glu	TGC Cys	TGC His	GGA Gly	GAT Asp	CTG Leu	CTT Leu	GAA Glu		828
			240				245					250				
TGT Cys	GCT Ala	GAT Asp	GAC Arg	AGG Arg	GCG Ala	GAC Asp	CTT Leu	GCC Ala	AAG Lys	TAT Tyr	ATC Ile	TGT Cys	GAA Glu	AAT Asn	CAA Gln	876
			255			260						265				
GAT Asp	TCG Ser	ATC Ile	TCC Ser	AGT Ser	AAA Lys	CTG Leu	AAG Lys	GAA Glu	TGC Cys	TGT Cys	GAA Glu	AAA Lys	CCT Pro	CTG Leu	TTG Leu	924
			270			275					280					
GAA Glu	AAA Lys	TCC Ser	CAC His	TGC Cys	ATT Ile	GCC Ala	GAA Glu	GTG Val	GAA Glu	AAT Asn	GAT Asp	GAG Glu	ATG Met	CCT Pro	GCT Ala	972
			285		290				295					300		
GAC Asp	TTG Leu	CCT Pro	TCA Ser	TTA Ala	GCT Ala	GCT Ala	GAT Asp	TTT Phe	GTT Val	GAA Glu	AGT Ser	AAG Lys	GAT Asp	GTT Val	TGC Cys	1020
				305				310					315			
AAA Lys	AAC Asn	TAT Tyr	GCT Ala	GAG Glu	GCA Ala	AAG Lys	GAT Asp	GTC Val	TTC Phe	CTG Leu	GGC Gly	ATG Met	TTT Phe	TTG Leu	TAT Tyr	1068



## (2) INFORMATION FOR SEQ ID NO: 2:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 585 amino acids  
 (B) TYPE: amino acid  
 (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

```

Asp Ala His Lys Ser Glu Val Ala His Arg Phe Lys Asp Leu Gly Glu
 1           5           10           15
Glu Asn Phe Lys Ala Leu Val Leu Ile Ala Phe Ala Gln Tyr Leu Gln
          20           25           30
Gln Cys Pro Phe Glu Asp His Val Lys Leu Val Asn Glu Val Thr Glu
          35           40           45
Phe Ala Lys Thr Cys Val Ala Asp Glu Ser Ala Glu Asn Cys Asp Lys
          50           55           60
Ser Leu His Thr Leu Phe Gly Asp Lys Leu Cys Thr Val Ala Thr Leu
          65           70           75           80
Arg Glu Thr Tyr Gly Glu Met Ala Asp Cys Cys Ala Lys Gln Glu Pro
          85           90           95
Glu Arg Asn Glu Cys Phe Leu Gln His Lys Asp Asp Asn Pro Asn Leu
          100          105          110
Pro Arg Leu Val Arg Pro Glu Val Asp Val Met Cys Thr Ala Phe His
          115          120          125
Asp Asn Glu Glu Thr Phe Leu Lys Lys Tyr Leu Tyr Glu Ile Ala Arg
          130          135          140
Arg His Pro Tyr Phe Tyr Ala Pro Glu Leu Leu Phe Phe Ala Lys Arg
          145          150          155          160
Tyr Lys Ala Ala Phe Thr Glu Cys Cys Gln Ala Ala Asp Lys Ala Ala
          165          170          175
Cys Leu Leu Pro Lys Leu Asp Glu Leu Arg Asp Glu Gly Lys Ala Ser
          180          185          190
Ser Ala Lys Gln Arg Leu Lys Cys Ala Ser Leu Gln Lys Phe Gly Glu
          195          200          205
Arg Ala Phe Lys Ala Trp Ala Val Ala Arg Leu Ser Gln Arg Phe Pro
          210          215          220
Lys Ala Glu Phe Ala Glu Val Ser Lys Leu Val Thr Asp Leu Thr Lys
          225          230          235          240
Val His Thr Glu Cys Cys His Gly Asp Leu Leu Glu Cys Ala Asp Asp
          245          250          255
Arg Ala Asp Leu Ala Lys Tyr Ile Cys Glu Asn Gln Asp Ser Ile Ser
          260          265          270
Ser Lys Leu Lys Glu Cys Cys Glu Lys Pro Leu Leu Glu Lys Ser His
          275          280          285
Cys Ile Ala Glu Val Glu Asn Asp Glu Met Pro Ala Asp Leu Pro Ser

```

290	295	300
Leu Ala Ala Asp Phe Val Glu Ser Lys Asp Val Cys Lys Asn Tyr Ala 305 310 315 320		
Glu Ala Lys Asp Val Phe Leu Gly Met Phe Leu Tyr Glu Tyr Ala Arg 325 330 335		
Arg His Pro Asp Tyr Ser Val Val Leu Leu Leu Arg Leu Ala Lys Thr 340 345 350		
Tyr Glu Thr Thr Leu Glu Lys Cys Cys Ala Ala Ala Asp Pro His Glu 355 360 365		
Cys Tyr Ala Lys Val Phe Asp Glu Phe Lys Pro Leu Val Glu Glu Pro 370 375 380		
Gln Asn Leu Ile Lys Gln Asn Cys Glu Leu Phe Glu Gln Leu Gly Glu 385 390 395 400		
Tyr Lys Phe Gln Asn Ala Leu Leu Val Arg Tyr Thr Lys Lys Val Pro 405 410 415		
Gln Val Ser Thr Pro Thr Leu Val Glu Val Ser Arg Asn Leu Gly Lys 420 425 430		
Val Gly Ser Lys Cys Cys Lys His Pro Glu Ala Lys Arg Met Pro Cys 435 440 445		
Ala Glu Asp Tyr Leu Ser Val Val Leu Asn Gln Leu Cys Val Leu His 450 455 460		
Glu Lys Thr Pro Val Ser Asp Arg Val Thr Lys Cys Cys Thr Glu Ser 465 470 475 480		
Leu Val Asn Arg Arg Pro Cys Phe Ser Ala Leu Glu Val Asp Glu Thr 485 490 495		
Tyr Val Pro Lys Glu Phe Asn Ala Glu Thr Phe Thr Phe His Ala Asp 500 505 510		
Ile Cys Thr Leu Ser Glu Lys Glu Arg Gln Ile Lys Lys Gln Thr Ala 515 520 525		
Leu Val Glu Leu Val Lys His Lys Pro Lys Ala Thr Lys Glu Gln Leu 530 535 540		
Lys Ala Val Met Asp Asp Phe Ala Ala Phe Val Glu Lys Cys Cys Lys 545 550 555 560		
Ala Asp Asp Lys Glu Thr Cys Phe Ala Glu Glu Gly Lys Lys Leu Val 565 570 575		
Ala Ala Ser Gln Ala Ala Leu Gly Leu 580 585		

## (2) INFORMATION FOR SEQ ID NO: 3:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 5 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: peptide



(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

Arg Tyr Thr Lys Lys  
1 5

(2) INFORMATION FOR SEQ ID NO: 4:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 5 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

Ala Tyr Thr Gln Gln  
1 5

(2) INFORMATION FOR SEQ ID NO: 5:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 39 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5:

CGTCAGACCT TGCATGCAGC CAAGACACCC TCACATAGC

39

(2) INFORMATION FOR SEQ ID NO: 6:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 42 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6:

CCGTTACGTT CTGTGGTGGC ATGCCCACTT CCAAGTCCAC CG

42

(2) INFORMATION FOR SEQ ID NO: 7:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 42 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 7:

CGCTCTCATA GTGGAARAGC TTCTAATAC GACAACTTCC CC

42

(2) INFORMATION FOR SEQ ID NO: 8:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 41 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 8:

CCCAAATGG TACCTGTGTC ATCACTCCTT GGGATAATAC C

41

(2) INFORMATION FOR SEQ ID NO: 9:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 44 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

## 33

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 9:

CCATCTGGAT CCAATGGTGC TTTGGCCAAA TAAATAGTTT CAGC

44

(2) INFORMATION FOR SEQ ID NO: 10:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 35 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 10:

GCCTCTTTTA CCGGTACAA GCTTGAGTCC ATTGG

35

(2) INFORMATION FOR SEQ ID NO: 11:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 37 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 11:

GGTAAGGTTT AGTCGACCTA TTTTGTGTTT TGTCTGC

37

(2) INFORMATION FOR SEQ ID NO: 12:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 44 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 12:

GGAAACGTAT GAATTCGATA TCATTGATAC AGACTCTGAG TACG

44

(2) INFORMATION FOR SEQ ID NO: 13:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 24 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(v) FRAGMENT TYPE: N-terminal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 13:

```
Met Leu Leu Gln Ala Phe Leu Phe Leu Leu Ala Gly Phe Ala Ala Lys
 1           5           10          15
Ile Ser Ala Asp Ala His Lys Ser
                20
```

(2) INFORMATION FOR SEQ ID NO: 14:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 4106 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(A) ORGANISM: *Saccharomyces cerevisiae*

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 14:

```
GAATTCTCTG TTGACTACTA AACTGAGAGA ATTTGCGAG ACTCTAAGAA CAGCTTTGAA      60
AGAGCGTTCT GCGGATGATT CCATAATTGT CACTCTGAGA GAGCAATGC AAAGAGAAAT      120
CTTCAGGTTG ATGTCGTTGT TCATGGACAT ACCTCCAGTG CAACCAACG AGCAATTCAC      180
TTGGGAATAC GTTGACAAG ACAAGAAAT CCACACTATC AAATCGACTC CGTTAGAATT      240
TGCTCCAAA TACGCAAAAT TGGACCCCTC CAGGCCACTC TCATTGATCA ATGATCCAAG      300
ACACCATATG GTAATTAAT TAAGATCGAT CGTTTAGGAA ACGTCCCTGG CGGAGATGCC      360
GTGATTACT TAAATGTTGA CAATGAARCA CTATCTAAAT TGGTTGTTAA GAGATTACAA      420
AATAACAAAG CTGCTTTTTT TGGATCTCAC ACTCCAAAGT TCATGGACAA GAAACTGGT      480
GTCATGGATA TTGAATTGTG GAACTATCCT GCCATGGCTA TAATTTACCT CAGCAAAAGG      540
CATCCGGTAT TAGATACCAT GAAAGTTTGA TGAATCATGC TATGTTGGAT CACTGGCTGC      600
CAGCTCGATG AAACCTCTAA ATTACCACTT CGCTACCGTC TGAAAATTC TGGGGTAAG      660
```

ACTCCGGTAA	AGACGGATTA	TACGTGATGA	CTCAAAAGTA	CTTCGAGGAG	TACTGCTTTC	720
AAATTGTGGT	CGATATCAAT	GAATTGCCAA	AAGAGCTGGC	TTCAAAATTC	ACCTCAGGTA	780
AGGAAGAGCC	GATTGTCTTG	CCCATCTGGA	CCCAATGGTG	CTTTGGCCAA	ATAAATAGTT	840
TCACGAGCTC	TGATGTAGAT	ACACGTATCT	CGACATGTTT	TATTTTACT	ATACATACAT	900
AAAAGAAATA	AAAATGATA	ACGTGTATAT	TATTATTTCAT	ATAATCAATG	AGGGTCATTT	960
TCTGAAACGC	AAAAAACGGT	AAATGAAAAA	AAAATAAAGA	TAGAAAAAGA	AAACAACAA	1020
AGGAAGGTTT	AGCATATTAA	ATAACTGAGC	TGATACTTCA	ACAGCATCGC	TGAAGAGAAC	1080
AGTATTGAAA	CCGAAACATT	TTCTAAGGC	AAACAAGGTA	CTCCATATTT	GCTGGACGTG	1140
TTCTTCTCT	CGTTTCATAT	GCATAATTCT	GTCAATAGCC	TGTTCTTTTT	CCTGGCTTAA	1200
ACATCCCGTT	TTGTAAGA	GAAATCTATT	CCACATATTT	CATTTCATCG	GCTACCATAC	1260
TAAGGATAAA	CTAATCCCGT	TGTTTTTGG	CCTCGTCACA	TAATTATAAA	CTACTAACCC	1320
ATTATCAGAT	GAAAGTGAGG	AAATATATTA	CTTTATGCTT	TTGGTGGGCC	TTTTCAACAT	1380
CCGCTCTTGT	ATCATCACAA	CAAATCCCAT	TGAAGGACCA	TACGTCACGA	CAGTATTTTG	1440
CTGTAGAAAG	CAATGAAACA	TTATCCCGCT	TGGAGGAAAT	GCATCCAAAT	TGGAATATG	1500
AACATGATGT	TCGAGGGCTA	CCAAACCATT	ATGTTTTTTC	AAAAGAGTTG	CTAAATTTGG	1560
GCAAAGATC	ATCATTAGAA	GAGTTACAGG	GGGATAACAA	CGACCACATA	TTATCTGTCC	1620
ATGATTIATT	CCCGGTAAC	GACCTATTTA	AGAGACTACC	GGTGCTGCT	CCACCAATGG	1680
ACTCAAGCTT	GTTACCGGTA	AAAGAAGCTG	AGGATAAACT	CAGCATAAAT	GATCCGCTTT	1740
TTGAGAGGCA	GTGGCACTTG	GTCAATCCAA	GTTTTCTTGG	CAGTGATATA	AATGTTCTTG	1800
ATCTGTGGTA	CAATAATATT	ACAGCGCGAG	GGGTCGTGGC	TGCCATTGTT	GATGATGGCC	1860
TTGACTACGA	AAATGAAGAC	TTGAAGGATA	ATTTTTCGCG	TGAAGGTTCT	TGGGATTTC	1920
ACGACAATAC	CAATTACCT	AAACCAAGAT	TATCTGATGA	CTACCATGGT	ACGAGATGTG	1980
CAGGTGAAT	AGCTGCCAAA	AAAGGTAACA	ATTTTTCGCG	TGTCGGGGTA	GGTTACAACG	2040
CTAAATCTC	AGGCATAAGA	ATCTTATCCG	GTGATATCAC	TACGGAAGAT	GAAGCTGCCT	2100
CCTTGATTTA	TGCTCTAGAC	GTAAACGATA	TATATTTCATG	CTCATGGGGT	CCCGCTGATG	2160
ACGGAAGACA	TTTACAAGGC	CCTAGTGACC	TGGTGAAAAA	GGCTTTAGTA	AAAGGTGTTA	2220
CTGAGGGAAG	AGATTCCAAA	GGAGCGATT	ACGTTTTTGC	CAGTGGAAAT	GGTGGAACTC	2280
GTGGTGATAA	TTGCAATTAC	GACGCTATA	CTAATTCCAT	ATATTCTATT	ACTATTGGGG	2340
CTATTGATCA	CAAAGATCTA	CATCCTCCTT	ATTCCGAAGG	TTGTTCCGCC	GTGATGGCAG	2400
TCACGTATTC	TTCAGGTTCA	GGCGAATATA	TTCAATTCGAG	TGATATCAAC	GGCAGATGCA	2460
GTATAGCCA	CGGTGGAAAC	TCTGCGGCTG	CTCCATTAGC	TGCCGGTGTT	TACACTTTGT	2520
TACTAGAGAC	CAACCCAAAC	CTAACTTGA	GAGACGTACA	GTATTATTATCA	ATCTTGCTCTG	2580
CGGTAGGGTT	AGAAAGAAC	GCTGACGGAG	ATTGGAGAGA	TAGCCCATG	GGGAGAAAT	2640

ACTCTCATCG CTATGGCTTT GGTAATATCG ATGCCCATAA GTTAATTGAA ATGTCCAAGA	2700
CCTGGGAGAA TGTTAACGCA CAACCTGGT TTTACCTGCC AACATTGTAT GTTCCCAGT	2760
CCACAAACTC CACGGAAGAG ACATTAGAAT CCGTCATAAC CATATCAGAA AAAAGTCTTC	2820
AAGATGCTAA CTTCAAGAGA ATTGAGCAGC TCACGGTAAC TGTAAGATAT GATACAGAAA	2880
TTAGGGGAAC TACGACTGTC GATTTAATAT CACCAGCGGG GATAATTCTA AACCTGGCG	2940
TTGTAAGACC AAGAGATGTT TCATCAGAGG GATTCAAAGA CTGGACATTC ATGCTGTAG	3000
CACATTGGGG TGAGAACGGC GTAGGTGATT GGAATCAAA GGTGAAGACA ACAGAAAATG	3060
GACACAGGAT TGACTTCCAC AGTTGGAGGC TGAAGCTCTT TGGGAATCC ATTGATTCAT	3120
CTAAACAGA AACTTTCGTC TTTGGAACG ATAAAGAGGA GGTGAACCA GCTGCTACAG	3180
AAAGTACCGT ATCACAATAT TCTGCCAGTT CACTTCTAT TTCCATCAGC GCTACTTCTA	3240
CATCTTCTAT CTCATTTGGT GTGGAACCGT CGGCCATTCC CCAACGACT ACTGCGAGTA	3300
CCGATCCTGA TTCTGATCCA AACACTCCTA AAAAATCTTC CTCTCCTAGG CAAGCCATGC	3360
ATTATTTTTT AACAAATATT TTGATTGGCG CCACATTTTT GGTGTTATAC TTCATGTTTT	3420
TTATGAATC AAGGAGAAGG ATCAGAAGGT CAAGAGCGGA AACGTATGAA TTCGATATCA	3480
TTGATACAGA CTCTGAGTAC GATTCTACTT TGGACAATGG AACTTCGGA ATTACTGAGC	3540
CCGAAGAGGT TGAGGACTTC GATTTTGATT TGCCGATGA AGACCATCTT GCAAGTTTGT	3600
CTTCATCAGA AAACGGTGAT GCTGAACATA CAATTGATAG TGTACTAACA AACGAAAATC	3660
CATTAGTGA CCCTATAAAG CAAAAGTTCC CAAATGACGC CAACGCAGAA TCTGCTTCCA	3720
ATAAATTACA AGAATTACAG CCTGATGTTC CTCCATCTTC CGGACGATCG TGATTCGATA	3780
TGTACAGAAA GCTTCAAATT ACAAAATAGC ATTTTTTCT TATAGATTAT AATACTCTCT	3840
CATACGTATA CGTATATGTC TATATGATAT ATAAACAAAC ATTAATATCC TATTCCTTCC	3900
GTTTGAATC CCTATGATGT ACTTTGCATT GTTGCACCC GCGAATAAAA TGAAACTCC	3960
GAACCGATAT ATCAAGCACA TAAAGGGGA GGTCCAATT AATGCATATT TAAGACCACA	4020
GCTGAATAAC TTTAAACGG CAGACAAAC AAAAATAGG TCGAATAAAC CTTACCTGCC	4080
TAGAAGCAAT GACAGCAGCT AATAAG	4106

## (2) INFORMATION FOR SEQ ID NO: 15:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 2526 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: DNA (genomic)

## (iii) HYPOTHETICAL: NO

## (iii) ANTI-SENSE: NO

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: *Saccharomyces cerevisiae*

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 15:

CCGTTTCTT	TTCGTA	AAAA	CAATAG	ACACTATATA	TAGACACTTT	TTCCTTTCCT	60
TCTTTGCGCG	ATTTCAAGAG	GAAAAGCATA	CTTAATAAG	AATATTCCTA	AAACACACGT		120
TCTGACGCGT	CAATTAGATC	GTCAGACCTT	GCATGCAGCC	AAGAACCCCT	CACATAGCAC		180
TGCCTCCTTC	CTCCTCTTTT	CTGTCAACAC	CTCACCTCCC	TCGTCCACTC	AACTGAGTGG		240
CTTTTCGCTC	CTTTTATACT	GCGCCATGAG	TAGTTTTCGT	TTCACGTATG	TGTCGAAAAA		300
AATTGAGGGT	TCATAAAAAA	ATTCGTGGAC	TTATTTATGG	AGAAACAGGG	AAATCCGACT		360
ACTTAAGAAA	AGGGTGTC	AGAGGATTTA	CTTTTTTCCT	TCTTTTGCA	TTTGTTCCCTA		420
TTTCCGCAAT	TGGACGGTTA	TTAAGAAGAA	CGCAATTGGC	TTTTCTGTAT	ATTAATAATAC		480
ATAGCGTAAT	AAAAAGATAA	GGTGAACACC	AAGCATATAG	TATAATATTA	CCTACCACAT		540
ATGAARCTGA	AACTGTAAAG	ATCTGCGGTC	CTTTCGTAC	TCTTTGCATC	GCAGGTTCTC		600
GGTAAGATAA	TACCAGCAGC	AAACAAGCGC	GACGACGACT	CGAATTC	CAAA	GTTCGTCAAG	660
TGCGCCCTTC	ATAAGCTTTA	CGGGGACTCG	CTAGAAAAATG	TGGGAAGCGA	CAAAAAACCG		720
GAAGTACGCC	TATTGAAGAG	GGCTGACGGT	TATGAAGAAA	TTATAATTAC	CAACCAGCAA		780
AGTTTCTATT	CGGTGGACTT	GGAAGTGGGC	ACGCCACCAC	AGAACGTAA	GGTCCTGGTG		840
GACACAGGCT	CCTCTGATCT	ATGGATTATG	GGCTCGGATA	ATCCATAC	CTG	TTCTTCGAAC	900
AGTATGGGTA	GTAGCCGGAG	ACGTGTTATT	GACAAACGTG	ATGATTCGTC	AAGCGGCGGA		960
TCTTTGATTA	ATGATATAAA	CCCATTGCG	TGGTTGACGG	GAACGGGCG	TGCCATTGGC		1020
CCCACTGCTA	CGGGCTTAGG	AGGCGGTTCA	GGTACGGCAA	CTCAATCCGT	GCCTGCTTCG		1080
GAAGCCACCA	TGACTGTCA	ACAATACGGG	ACATTTTCCA	CTTCGGGCTC	TTCTACATTT		1140
AGATCAAACA	ACACCTATTT	CAGTATTAGC	TACGGTGATG	GGACTTTTGC	CTCCGGTACT		1200
TTTGATACGG	ATGTTTGGGA	TTTAAGCGAC	TTGAACGTTA	CCGGGTGTC	TTTTGCCGTT		1260
GCCATGAAA	CGAATCTTAC	TATGGGTGTG	TTAGGTATTG	GTTCGCCCGA	ATTAGAAGTC		1320
ACTTATCTCG	GCTCTACTGC	GTCTCATAGT	GGAAGAGCTT	ATAAATACGA	CAACTTCCCC		1380
ATTGTATTGA	AAATTTCTGG	TGCTATCAAA	AGCAACACAT	ATTCTTTGTA	TTTGAACGAC		1440
TCGGACGCTA	TGCATGGCAC	CATTTTGTTT	GGAGCCGTGG	ACCACAGTAA	ATATAACGGC		1500
ACCTTATACA	CAATCCCAT	CGTAACACT	CTGAGTGCTA	GTGGATTTAG	CTCTCCCAT		1560
CAATTGTATG	TCACTATTAA	TGGTATCGGT	ATTAGTGATT	CTGGGAGTAG	TAAACAGACC		1620
TTGACTACCA	CTAAAATACC	TGCTTTGTCG	GATTCCGGTA	CTACTTTGAC	TTATTTACCT		1680
CAAACAGTGG	TAGTATGAT	CGCTACTGAA	CTAGGTGCGC	AATACTCTTC	CAGGATAGGG		1740
TATTAGGTAT	TGACTGTCC	ATCTGATGAT	AGTATGGAAA	TAGTGTTCGA	TTTTGGTGGT		1800
TTTCACATCA	ATGCACCACT	TTGAGTTTTT	ATCTTGAGTA	CTGGCACTAC	ATGCTTTTTA		1860
GGTATTATCC	CAACGAGTGA	TGACACAGGT	ACCATTTTGG	GTGATTCA	TTTGACTAAC		1920

GCGTACGTGG TTTATGATTT GGAGAATCTT GAAATATCCA TGGCACAAGC TCGCTATAAT	1980
ACCACAAGCG AAAATATCGA AATTATCACA TCCTCTGTTC CAAGCGCCGT AAAGGCACCA	2040
GGCTATACAA ACACTTGGTC' CACAAGTGCA TCTATTGTTA CCGGTGGTAA CATATTTACT	2100
GTAATTCCT CACAAACTGC TTCCTTTAGC GGTAACTGA CGACCACTAC TGCATCCGCC	2160
ACTTCTACAT CAAGTAAAG AATGTTGGT GATCATATAG TTCCATCTTT ACCCCTCACA	2220
TTAATTTCTC TTCTTTTTCG ATTCATCTGA AAACCGTTGC ACAAAGTTTA GACATTCACA	2280
TCTCCAAGCG AGTTGGAGTT TCTGGCGGAA ATCGTTGCTC TCGCTTGGGC AAAGTTTTTT	2340
TTTATTATTA ATTTTTTATT GTTACGTTGG CGGTCITTAT TTTTACTTCA CAATAGTTTA	2400
TCTTACCCAC TAAGAATAGG TTACCATTTA TTCACATTTT TTTTCTCAT TCCTAGTATA	2460
CTATTTACCT GGGATATGGC CTATAATCAA AGGCTTTAAT ATTCTAATAA TTCGTTTGGC	2520
ATCTAG	2526




CLAIMS

1. A process for preparing albumin by secretion from a yeast genetically modified to produce and secrete the albumin, comprising culturing the yeast in a culture medium such that albumin is secreted into the medium, characterised in that the yeast cells have a reduced level of yeast aspartyl protease 3 proteolytic activity.
2. A process according to Claim 1 wherein the said proteolytic activity is an endoprotease activity specific for monobasic sites and for paired basic amino acids in a polypeptide.
3. A process according to Claim 1 or 2 wherein the yeast is *S. cerevisiae*.
4. A process according to Claim 1, 2 or 3 wherein the yeast lacks a functional *YAP3* gene or homologue thereof.
5. A process according to any one of Claims 1 to 4 wherein the yeast cells additionally have a reduced level of *S. cerevisiae* Kex2p proteolytic activity.
6. A process according to any one of the preceding claims wherein the albumin is a human albumin.
7. A culture of yeast cells containing a polynucleotide sequence encoding an albumin and a second polynucleotide sequence encoding a secretion signal causing albumin expressed from the first polynucleotide sequence to be secreted from the yeast, characterised in that the yeast cells have a reduced level of yeast aspartyl protease 3 proteolytic


activity.

8. A culture according to Claim 7 wherein the albumin is a human albumin.
- 5 9. A culture according to Claim 7 or 8 wherein the yeast is *S. cerevisiae*.
- 10 10. A culture according to any one of Claims 7 to 9 wherein the said signal is cleaved by the yeast prior to release of the albumin from the yeast.
- 15 11. A culture according to any one of Claims 7 to 10 wherein the yeast cells additionally have a reduced level of Kex2p proteolytic activity.
12. A culture according to Claim 11 wherein the said secretion signal is cleaved from the albumin by a protease other than Kex2p.
- 20 13. A modified albumin having at least 90% sequence identity to a naturally-occurring albumin, which naturally-occurring albumin is susceptible to cleavage with yeast aspartyl protease 3 (Yap3p) when expressed and secreted in yeast, characterised in that the modified albumin is not susceptible to such cleavage.
- 25 14. A modified albumin according to Claim 13 wherein the modified albumin lacks a monobasic amino acid present in the naturally-occurring albumin protein.
- 30 15. A modified albumin according to Claim 13 or 14 wherein the said monobasic amino acid is arginine.

16. A modified albumin according to Claim 14 or 15 wherein the modified albumin additionally lacks a pair of basic amino acids present in the naturally-occurring albumin.
- 5 17. A modified albumin according to Claim 16 wherein the said pair of amino acids is Lys, Lys; Lys, Arg; Arg, Lys; or Arg, Arg.
18. A modified albumin according to Claim 13 wherein the naturally-occurring albumin is a human albumin and the modified protein lacks  
10 Arg<sup>410</sup>; and, optionally, residues 413 and 414 do not each consist of lysine or arginine.
19. A modified albumin according to Claim 18 which is a human albumin having the amino acid changes R410A, K413Q, K414Q.  
15
20. A polynucleotide encoding a modified albumin according to any one of Claims 13 to 19.
21. A yeast containing a polynucleotide according to Claim 20,  
20 transcription signals such that the modified albumin is expressed in the yeast, and a further polynucleotide adjacent the said polynucleotide such that the modified albumin is secreted from the yeast.

1/5  
**Figure 1**


2/5  
**Figure 2**





3/5


Figure 3

5' and 3' regions of YAP3 obtained by PCR:




4/5


Figure 4Figure 5

5/5  
Figure 65' and 3' regions of *KEX2* obtained by PCR:



Cloned into pUC19

*Bam*HI/*Hind*III  
fragment  
isolated

Cloned into pUC19

*Hind*III/*Sal*I  
fragment  
isolated

Subcloned into pUC19HX

*LYS2* *Hind*III  
fragment inserted



## INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 95/00434

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N15/12 C12N1/21 A61K38/38 C12P21/02 C12N15/11  
C12N1/20

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N A61K C12P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, A	BIOCHIMIE, vol. 76, 1994 pages 226-233, Y. BOURBONNAIS ET AL. 'Cleavage of prosomatostatins by the yeast Yap3 and Kex2 endoprotease' *see the whole article* ---	1-21
Y	BIOTECHNOLOGY, vol. 8, no. 1, 1990 pages 42-46, D. SLEEP ET AL. 'The secretion of human serum albumin from the yeast <i>S. cerevisiae</i> using five different leader sequences' *see the whole article* ---	1-21
	--- -/--	

☒ Further documents are listed in the continuation of box C.☐ Patent family members are listed in annex.

## \* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance  
 \*E\* earlier document but published on or after the international filing date  
 \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)  
 \*O\* document referring to an oral disclosure, use, exhibition or other means  
 \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention  
 \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone  
 \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art  
 \*&\* document member of the same patent family

Date of the actual completion of the international search

23 June 1995

Date of mailing of the international search report

- 5. 07. 95

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
 NL - 2280 HV Rijswijk  
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
 Fax (+31-70) 340-3016

Authorized officer

Marie, A

## INTERNATIONAL SEARCH REPORT

Int ional Application No  
PCT/GB 95/00434

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	EMBO JOURNAL, vol. 12, no. 1, 1993 pages 285-294, Y. BOURBONNAIS ET AL. 'Isolation and characterisation of S. cerevisiae mutants defective in somatostatin expression...' *see the whole article* ---	1-21
Y	YEAST, vol. 6, 1990 pages 127-137, M. EGEL-MITANI ET AL. 'A novel aspartyl protease allowing KEX2-independent MF alfa propheromone processing in yeast' *see the whole article* -----	1-21